



P.O. Box 8516
Tarrytown, New York 10591-8516
914.372.2331

THE TRUTH: MALE CIRCUMCISION DOES NOT PREVENT HIV

Campaigns Promoting Male Circumcision Are a Misleading and Dangerous Mistake

Summary

Mass male circumcision is being promoted as a method of curbing the AIDS pandemic in sub-Saharan Africa. Stopping the spread of HIV requires using available resources strategically, and circumcision's costs and harms are too significant to ignore. Mass circumcision campaigns will divert resources from proven prevention programs, result in a high number of complications, increase risk-compensation behaviors, and put women at higher risk for HIV.

Circumcision is an expensive and risky procedure that was shown to reduce risk by 50–60% for heterosexual males only in three highly controlled, short-term clinical trials. However, condom promotion and safe-sex education have already been shown to reduce infection rates more effectively for both males and females, at a lower cost. Furthermore, anti-retroviral drugs have shown a promising 92% reduction in HIV transmission.¹

Adult males are vulnerable to the belief that circumcision offers them immunity from HIV,² raising ethical concerns about promoting adult male circumcision, and questions regarding the effectiveness of the intervention.

Some have proposed circumcising infants, but this, too, has ethical ramifications.³ Removing healthy tissue from children deprives them of their right to autonomy. Surgery of any kind places them at immediate risk from complications, while the HIV benefit, if any, is 15–20 years away.

Male circumcision does not protect women;⁴ in fact, it may increase their risk of contracting HIV.⁵ Further, circumcision does not protect men who have sex with men.^{6 7}

Background

Results from three randomized clinical trials (RCTs) in Africa, showing a reduction in female-to-male transmission of HIV after circumcision, have resulted in the promotion of mass male circumcision to reduce HIV transmission.^{8 9 10} This has been followed by the mobilization of vast sums of money from international health agencies and foundations to roll out mass surgical interventions in sub-Saharan Africa, and to influence male infant circumcision policy in the United States.

RCTs not applicable to the Real World HIV Battle

A number of factors warrant caution in extrapolating the RCT results to larger populations. Unlike in a real-world setting, study participants were provided free condoms, extensive education and counseling, and they were paid, making comparisons to the general population questionable at best. The only certain conclusion that can be drawn from the RCTs is that while circumcision might *delay* HIV infection for circumcised males at high risk, it has no effect on infection rates for women, and is irrelevant for children.

There is insufficient evidence that circumcision as an HIV-prevention strategy is efficacious or would ultimately save lives. A 50–60% transmission reduction in a population engaging in high-risk behaviors is ultimately not effective, especially when it promotes a false sense of immunity.

Two recent studies examining African circumcision rates and HIV prevalence found that circumcision status was not significantly associated with HIV. One study examined data from 13 sub-Saharan countries found no association,¹¹ and another found that circumcision made no difference in HIV rates in South Africa.¹² A 2007 study concluded that, once commercial sex-worker patterns are factored in, male circumcision is not significantly associated with lower HIV.¹³

Another 2009 publication also shows that circumcision status is not correlated with lower HIV prevalence rate—national household survey data from 18 countries in which circumcision status was tracked showed that HIV was higher in circumcised males for 10 of the 18 countries (Cameroon, Guinea, Haiti, Lesotho, Malawi, Niger, Rwanda, Senegal, Tanzania, and Zimbabwe).¹⁴

Recent evidence demonstrates that Langerhans cells in the foreskin have a protective effect against pathogens, including HIV, by secreting langerin.¹⁵ The original theory (which led to promoting circumcision to stop HIV) was that Langerhans cells are an entry point for viruses. It now appears that the theory was partially true, but that the mechanism at work is that Langerhans cells set a trap for viruses in order to destroy them with langerin. It is plausible that Langerhans cells are only infected by HIV when they are challenged by a high viral load.

Male Circumcision May Increase Risk of HIV

The long-term consequences of promoting circumcision could actually worsen the HIV epidemic by promoting a false sense of security and undermining safe-sex practices and condom usage.^{16 17} African men are already lining up to be circumcised, believing they will no longer need to use condoms.^{18 19 20} A 2009 South African National Communication Survey on HIV/AIDS found that 15% of men and women held the mistaken belief that circumcision makes sex without condoms safe.²¹

Assuming the 50–60% protective effect found in the RCTs is valid, and if all African males were circumcised over the next 15 years and all other conditions were held constant, circumcision would reduce the number of infections by 8%, and related deaths by 1%.²² It is unknown how many additional infections and deaths would occur from HIV-positive circumcised men transmitting the virus to their female partners, and from the iatrogenic spread of the virus—not to mention the number of circumcision complications that would occur.

HIV infections have been found following the circumcision of virgins, both male and female, indicating that circumcision spreads iatrogenically—probably from multiple use of contaminated instruments.^{23 24} Also troubling as an indicator of what is to come is the recent announcement that in Swaziland, circumcised males are being targeted as blood donors, because they are considered to be “the safe group for blood donation”²⁵—irrespective of whether the individuals have been tested for HIV. In resource-poor countries of the world, blood transfusion remains a major avenue of transmission of HIV, and much of the blood supply is not tested.²⁶

Male Circumcision Endangers Women

Male circumcision not only offers no protection to women,²⁷ but it increases the risk to women if sex is resumed before the wound has completely healed.²⁸ A recent WHO study found that one-quarter of circumcised males still had not healed sixty days after the surgery.²⁹ Further, circumcision places women at greater risk of unsafe sex practices if they or their circumcised male partners wrongly believe they are immune to HIV.

Circumcision Will Result in Burdensome Complications

A recent issue of the WHO Bulletin noted that African ritual circumcisions have a 35% complication rate, while clinical circumcisions have an 18% complication rate.²⁹ A neonatal circumcision complication rate of 20.2% was found in Nigeria.³⁰ Dealing with these complications will divert even more resources away from other programs.

Unethical Medical Practice

Circumcision permanently removes healthy, functional, beneficial tissue.³¹ Mass prophylactic surgery is unprecedented, and there are sensible, safe and effective alternatives.³² Circumcised men still must wear condoms for protection (as well as to protect their sexual partners), and there is no evidence that being circumcised and wearing condoms is any better than wearing condoms alone. Further, because circumcision can actually increase the risk of infection with HIV, and in view of the complications and negative consequences of the surgery, promoting it as a public health measure is unethical.

Informed Consent

For fully informed consent to occur, men must be educated about the risks and sensory losses from circumcision, as well as be made aware that it does not offer full protection and that they will still need to wear condoms during sex. The number of reports emerging regarding African males agreeing to circumcision so they will no longer need to use condoms reveals that fully informed consent is not always occurring.^{33 34 35}

Effective Prevention Programs Already Exists

Education, safe-sex practices, and consistent condom use are proven, effective measures of curbing HIV transmission. Uganda demonstrated a 47% reduction in HIV prevalence from increased safe-sex education and condom promotion; this social-prevention program is available now, is highly effective, and does not involve the numerous risks and losses from surgery.³⁶ A 2008 study revealed that condoms are 98% effective at hindering HIV transmission, and 95 times more cost-effective than circumcision.³⁷ Consistent condom use reduces lifetime risk by 20%³⁸ as compared to circumcision's 8%.²⁴ A recent report from South Africa shows condom use significantly increased from 2002 to 2008, and the HIV rates finally began to level off.³⁹ Diverting resources to circumcision will impede this progress.

A study using mathematical impact modeling of circumcision, anti-retroviral therapy, and condom use in South Africa concluded: "Male circumcision was found to have considerably lower impact than condom use or antiretroviral therapy on HIV infection and death rates."⁴⁰ Another study showed that anti-retroviral drugs reduced transmission by 92%, far more effective than any reduction claimed by proponents of circumcision.

Conclusion

Promoting an intervention that at best reduces the probability of infection for males only, while creating a false sense of security and risk of iatrogenic HIV transmission, is irresponsible and unethical. Circumcision offers no protection for men who have sex with men, and may increase the risk to women. Male circumcision will result in unacceptable complications, the treatment of which will further burden the inadequate healthcare infrastructure of developing countries. Promoting male circumcision drains resources that should be devoted to proven measures such as condom promotion, increased safe-sex education, and increased anti-retroviral drug distribution.

References

- ¹ Available online at: http://www.newsmxhealth.com/health_stories/drugs_slash_HIV_risk/2010/05/28/315637.html
- ² Gusongoirye D. Rwanda: Nothing can fight HIV/AIDS better than discipline. *The New Times (Kigali)* February 12. (2008)
- ³ Somerville M. Altering baby boys' bodies: The ethics of infant male circumcision. In: *The Ethical Canary: Science, Society, and the Human Spirit*. New York: Viking, 2000.
- ⁴ Turner, et al. Men's circumcision status and women's risk of HIV acquisition in Zimbabwe and Uganda. *AIDS* 2007, 21:1779-1789.
- ⁵ Wawer M, Kigozi G, Serwadda D, et al. Trial of male circumcision in HIV+ men, Rakai, Uganda: effects in HIV+ men and in women partners. 15th Conference on Retroviruses and Opportunistic Infections. 3-6 February 2008. Boston. Abstract 33LB.
- ⁶ Templeton D. J., Jin F., Prestage G. P., et al. Circumcision status and risk of HIV seroconversion in the HIM cohort of homosexual men in Sydney. In 4th Conference on the HIV Pathogenesis, Treatment and Prevention. 23-25 July 2007. Sydney, Australia: International AIDS Society.
- ⁷ Millett G, Ding H, Lauby J, Flores S, Stueve A, Bingham T, et al. Circumcision Status and HIV Infection Among Black and Latino Men Who Have Sex With Men in 3 US Cities. *J Acquir Immune Defic Syndr*. 2007;46(5):643-650.
- ⁸ Auvert B, Taljaard D, Lagarde E, Sobngwi-Tambekou J, Sitta R, Puren A. Randomized, controlled intervention trial of male circumcision for reduction of HIV infection risk: the ANRS 1265 trial. *PLoS Med* 2005;2(11): e2.
- ⁹ Bailey RC, Moses S, Parker CB, Agot K, Maclean I, Krieger JN, et al. Male circumcision for HIV prevention in young men in Kisumu, Kenya: A randomised controlled trial. *Lancet*. 2007;369(9562):643-56.
- ¹⁰ Gray RH, Kigozi G, Serwadda D, Makumbi F, Watya S, Nalugoda F, et al. Male circumcision for HIV prevention in men in Rakai, Uganda: A randomised trial. *Lancet*. 2007;369(9562):657-66.
- ¹¹ Michel Garenne. *African Journal of AIDS Research* 2008, 7(1): 1-8.
- ¹² Connolly C, et al. Male circumcision and its relationship to HIV infection in South Africa: Results of a national survey in 2002. *S Afr Med J* 2008;98:789-94.
- ¹³ Talbot JR. Size Matters: The number of prostitutes and the global HIV/AIDS pandemic. *PLoS One*. 2007;2(6): e543.
- ¹⁴ Mishra, Vinod, Amy Medley, Rathavuth Hong, Yuan Gu, and Bryant Robey. 2009. Levels and Spread of HIV Seroprevalence and Associated Factors: Evidence from National Household Surveys. DHS Comparative Reports No. 22. Calverton, Maryland, USA: Macro International Inc. p. 209.
- ¹⁵ de Witte L, Nabatov A, Pion M, Fluitsma D, de Jong, MAWP, de Gruijl, T, et al. Langerin is a natural barrier to HIV-1 transmission by Langerhans cells. *Nat Med*. 2007;(3):367-71.
- ¹⁶ Kalichman S., Eaton L., Pinkerton S. Circumcision for HIV prevention: failure to account for behavioral risk compensation *PLoS Med*. 2007;4(3):e137-138.
- ¹⁷ Myers A., Myers J. Male circumcision-the new hope? *S Afr Med J*. 2007;97(5):338-341.
- ¹⁸ Gusongoirye D. Rwanda: Nothing can fight HIV/AIDS better than discipline. *The New Times (Kigali)* February 12, 2008.
- ¹⁹ Nyakairu, F. Uganda turns to mass circumcision in AIDS fight. *Reuters Africa*. 13 August, 2008.
- ²⁰ IRIN, Swaziland: Circumcision gives men an excuse not to use condoms. *Humanitarian news and analysis UN Office for the Coordination of Humanitarian Affairs*. 31 July, 2008.
- ²¹ South African Government. South African National Communication Survey on HIV/AIDS, 2009. http://www.info.gov.za/issues/hiv/survey_2009.htm.
- ²² Williams BG, Lloyd-Smith JO, Gouws E, et al. The potential impact of male circumcision on HIV in Sub-Saharan Africa. *PLoS Med* 2006;3:e262.
- ²³ Brewer DD, Potterat JJ, Roberts Jr, JM, et al. Male and female circumcision associated with prevalent HIV infection in virgins and adolescents in Kenya, Lesotho, and Tanzania. *Ann Epidemiol*. 2007;17:217-226.
- ²⁴ Stallings RY, Karugendo E. Female circumcision and HIV infection in Tanzania: for better or for worse? Third International AIDS Society Conference on HIV Pathogenesis and Treatment. Rio de Janeiro, 25-27 July, 2005.
- ²⁵ Masilela C. Circumcised can still donate blood -SNBTS. *The Swazi Observer*, 17 June, 2010. Available online at: <http://www.observer.org.sz/index.php?news=14149>
- ²⁶ Chamberland ME, Lackritz EM, Busch MP. HIV screening of the blood supply in developed and developing countries. *AIDS Rev* 2001;3:24-35.
- ²⁷ Turner, et al. Men's circumcision status and women's risk of HIV acquisition in Zimbabwe and Uganda. *AIDS* 2007, 21:1779-1789.
- ²⁸ Wawer M, Kigozi G, Serwadda D, et al. Trial of male circumcision in HIV+ men, Rakai, Uganda: effects in HIV+ men and in women partners. 15th Conference on Retroviruses and Opportunistic Infections. 3-6 February 2008. Boston. Abstract 33LB.
- ²⁹ Bailey RC, Egesah O, Rosenberg S. Male circumcision for HIV prevention: a prospective study of complications in clinical and traditional settings in Bungoma, Kenya. *Bull WHO*. 2008;86(9): 669-677.
- ³⁰ Okeke LI, Asinobi AA, Ikerowo OS. Epidemiology of complications of male circumcision in Ibadan, Nigeria. *BMC Urology*. 2006;6:21.
- ³¹ Cold C.J., Taylor J. R. The prepuce. *BJU Int*. 1999;83Suppl.1:34-44.
- ³² Fox M, Thomson M. Short changed: the law and ethics of male circumcision. *Int J Children's Rights*. 2005;13:161-181.
- ³³ IRIN, Swaziland: Circumcision gives men an excuse not to use condoms. *UN Office for the Coordination of Humanitarian Affairs*. 31 July, 2008. Available online at: <http://www.irinnews.org/Report.aspx?ReportId=79557>
- ³⁴ Nyakairu, F. Uganda turns to mass circumcision in AIDS fight. *Reuters Africa*. 13 August, 2008. Available online at: <http://africa.reuters.com/top/news/usnBAN338078.html>
- ³⁵ Gusongoirye D. Rwanda: Nothing can fight HIV/AIDS better than discipline. *The New Times (Kigali)* February 12, 2008.
- ³⁶ Low-Beer D, Stoneburner RL. Behaviour and communication change in reducing HIV: Is Uganda unique? *Johannesburg: Centre For Aids Development, Research And Evaluation*. 2004, 14 p.
- ³⁷ McAllister RG, Travis JW, Bollinger D, Rutiser C, Sundar V: The Cost to Circumcise Africa. *Int. J. Men's Health* 2008;7(2):307-316.
- ³⁸ Hallett TB, Gregson S, Lewis JJ, Lopman BA, Gamett GP. Behaviour change in generalised HIV epidemics: impact of reducing cross-generational sex and delaying age at sexual debut. *Sex Transm Infect* 83 2007;(suppl 1): i50-i54.
- ³⁹ Available online at: <http://news.bbc.co.uk/go/pr/fr/-/1/hi/world/africa/8091489.stm>
- ⁴⁰ Lima V, et al. The combined impact of male circumcision, condom use and HAART coverage on the HIV-1 epidemic in South Africa: a mathematical model. 5th IAS Conference on HIV Treatment, Pathogenesis and Prevention, Cape Town, abstract WECA105, 2009.